



Study of Serum Uric Acid in Acute Myocardial Infarction

Dr.Shilpa.R¹, Dr.SaiKrishna. K²

¹Assistant Professor in Dept of General Medicine, Dr.PSIMS& RF, Gannavaram, Vijayawada, India.

²Consultant in Medical Oncology, American Oncology Institute, Vijayawada, India.

Corresponding Author: Dr.Shilpa.R

ABSTRACT:

BACKGROUND:

Acute myocardial infarction (AMI) is overwhelmingly the most important form of ischemic heart disease (IHD) which continues to be the leading cause of death in the industrialized and developing countries like India, despite spectacular progress in their prevention, detection and treatment over the last three decades. The relation between uric acid and cardiovascular disease is observed not only with frank hyperuricemia (defined as more than 6mg/dl in women & more than 7 mg/dl in men) but also with uric acid levels considered to be normal but at high range.

AIMS AND OBJECTIVES: To study the serum uric acid level of patients of Acute Myocardial Infarction admitted to our hospital. To correlate the serum uric acid levels on day 0,3,7 and comparing with killips class with coefficient of correlation and to assess the morbidity and mortality of these patients.

METHODS: This hospital based case study was performed in Dr.PSIMS& RF, Chinoutpalli, Krishna district, Andhra Pradesh, south India from October 2016 to September 2017. A total of 75 cases of Acute MI were studied. All the subjects were interviewed, examined and investigated as per the predesigned proforma.

RESULTS: 75 cases of acute MI were included in the study. Mean SUA for discharged patients was 4.67 ± 1.95 /dl and it was 7.1 ± 1.45 mg/dl for the patients who died in the hospital. SUA levels were significantly higher in the patients who succumbed as compared to those who were discharged from the hospital ($p = 0.000$).

CONCLUSION:

Serum uric levels are correlated with killip's classification. Patients of higher killips class have higher uric acid levels. Combination of killip's class and serum uric acid levels after acute myocardial infarction is a good predictor of mortality. Serum uric acid levels were high in the patients who died in the seven day follow up period and the patients who died were also in higher killip's class.

KEYWORDS: Acute Myocardial Infarction (AMI); Serum Uric acid (SUA); Killip's class; Framingham study

Received 14 Nov., 2024; Revised 25 Nov., 2024; Accepted 27 Nov., 2024 © The author(s) 2024.

Published with open access at www.questjournals.org

I. INTRODUCTION:

Acute myocardial infarction (AMI) is the most important form of ischemic heart disease (IHD) which continues to be the leading cause of death in the industrialized and developing countries like India¹ A large number of asymptomatic individuals are at serious risk of developing myocardial infarction (MI) because of their genetic predisposition, smoking behaviour and sedentary lifestyle. Epidemiological studies have recently shown that uric acid may be a risk factor for cardiovascular diseases and a negative prognostic marker for mortality in subjects with pre-existing heart failure. Many studies including the National Health and Nutrition Examination Survey (NHANES) study 2 concluded that uric acid is an independent risk factor for development of cardiovascular and cerebrovascular diseases. Framingham Heart study³ concluded that an association between hyperuricemia and cardiovascular diseases merely reflects the link between serum uric acid and other risk factors, including hypertension, renal disease, elevated lipoprotein levels and the use of diuretics. Adenosine synthesized locally by vascular smooth muscle in cardiac tissue is rapidly degraded by the endothelium to uric acid, which undergoes rapid efflux to the vascular lumen due to low intracellular pH and negative membrane potential⁷ Xanthine oxidase⁸ and uric acid⁹ synthesis are increased in vivo under ischemic conditions, and therefore elevated serum uric acid may act as a marker of underlying tissue ischemia. Although the mechanism

by which uric acid may play a pathogenetic role in cardiovascular disease is unclear, hyperuricemia is associated with deleterious effects on endothelial dysfunction, oxidative metabolism, platelet adhesiveness, and aggregation. Uric acid may have direct role in atherosclerotic process because atherosclerotic plaque contains more uric acid than control arteries. Hyperuricemia via purine metabolism may also promote thrombus formation.⁴ The relation between uric acid and cardiovascular disease is observed not only with frank hyperuricemia (defined as more than 6mg/dl in women & more than 7 mg/dl in men) but also with uric acid levels considered to be normal but at high range.⁵This study was done to determine whether raised serum uric acid levels were an independent risk factor for acute MI and to determine its prognostic importance if any.

II. MATERIALS AND METHODS:


SOURCE OF DATA COLLECTION: The study was conducted at Dr.PSIMS& RF, Chinoutpalli , Krishna district, Andhra Pradesh , from October 2016 to September 2017. A total of 75cases of Acute MI were studied. All the subjects were interviewed, examined and investigated as per the predesigned proforma.Patients more than 18 years of age diagnosed to have acute MI who presented to hospital within 24 hours of onset of symptoms were included in the study.Acute MI was defined as, ‘increased myocardial enzyme concentrations with typical chest pain persisting more than 30 minutes or electrocardiographic changes (including ischemic ST-segment depression, ST-segment elevation or pathologic Q waves). Increased enzyme concentrations were defined as peak creatine phosphokinase level more than 2 times upper limit of normal.⁶Patients with a condition known to elevate uric acid level e.g. chronic kidney disease, gout, haematological malignancy and hypothyroidism were excluded.Patients receiving drugs affecting serum UA levels (diuretics, ethambutol, pyrizinamide, salicylates, losartan, allopurinol, probenecid, atorvastatin, fenofibrate).Chronic alcoholics.

III. RESULTS:

Table 1
URIC ACID LEVELS DAY 0

KILLIPS CLASS	<4.1	4.1-5.5	5.5-7	>7	TOTAL
1	21	21	3	2	47
2	4	1	7	0	12
3	1	0	8	3	12
4	0	0	0	4	4
TOTAL	26	22	18	9	75

Graph 1:



CROSS TABULATION BETWEEN SERUM URIC ACID AND KILLIPS CLASS ON DAY OF ADMISSION


On applying chi-square test ,p value significant (<0.05)As p value is significant,so there is association between serum uric acid concentration and killips class on day of admission. serum uric acid levels are higher in patients who are in higher killips class.

URIC ACID LEVELS DAY 3

Table 2

KILLIPS CLASS	<4.1	4.1-5.5	5.5-7	>7	TOTAL
1	28	13	6	0	47
2	1	1	3	0	5
3	1	1	5	6	13
4	0	0	0	9	9
TOTAL	30	15	14	15	74

Graph 2:



CROSS TABULATION BETWEEN SERUM URIC ACID AND KILLIPS CLASS ON DAY 3


On applying chi-square test ,p value significant (<0.05) As p value is significant,so there is association between serum uric acid concentration and killips class on day of 3,serum uric acid levels are higher in patients who are in higher killips class.

URIC ACID LEVELS DAY 7

Table 3:

KILLIPS CLASS	<4.1	4.1-5.5	5.5-7	>7	TOTAL
1	31	11	10	1	53
2	0	0	6	0	6
3	0	0	1	6	7
4	0	0	0	3	3
TOTAL	31	11	17	10	69


Graph 3:



CROSS TABULATION BETWEEN SERUM URIC ACID AND KILLIPS CLASS ON DAY7.

On applying chi-square test, p value significant (<0.05) As p value is significant, so there is association between serum uric acid concentration and Killips class on day 7. Serum uric acid levels are higher in patients who are in higher Killips class.

Graph 4:



FREQUENCY DISTRIBUTION OF PATIENTS ACCORDING TO KILLIPS CLASS ON DAY 0, DAY 03, DAY 07.

Table 4:

KILLIPS CLASS	DAY 0 N(%)	DAY 03 N(%)	DAY 07 N(%)
1	45(60%)	47(64%)	52(75%)
2	11(15%)	6 (8%)	9 (14%)
3	14(19%)	9 (13%)	6 (8%)
4	5(6%)	9 (15%)	2 (3%)
TOTAL	75	74	69

From table 4, it is observed that patients have shifted from class 2 and 3 to class 1 and 4. one patient expired till day 3, and 5 till day 7, total 6 patients died during 7 day follow up.

Table 5: RELATION BETWEEN SERUM URIC ACID AND MI OUTCOME

Outcome	Discharge (n=69)	Death (n=6)	p value
Mean SUA ± SD (in mg/dl)	4.67 ±1.95	7.1 ±1.45	0.00
Range (in mg/dl)	3.2-7.8	5.8-9.6	

Mean SUA for discharged patients was 4.67 ±1.95 /dl and it was 7.1 ±1.45 mg/dl for the patients who died in the hospital. SUA levels were significantly higher in the patients who succumbed as compared to those who were discharged from the hospital (p = 0.000).

IV. DISCUSSION:

The well recognized risk factors like age, male sex, smoking, diabetes, hypertension, metabolic syndrome explain only a part of the MI cases. Hence a search for other risk factors is the need of the hour. Many studies have found conflicting role of uric acid in patients with cardiovascular disease.

It has been reported that UA promotes vascular smooth muscle proliferation and upregulates the expression of platelet-derived growth factor and monocyte chemoattractant protein-1⁷. Hypoxanthine is converted to uric acid via xanthine. This reaction can be catalyzed by xanthine hydrogenase and xanthine oxidase, the latter of which produces uric acid and superoxide. Thus, it is possible that, in certain diseased conditions, hyperuricemia is accompanied by the increased production of reactive oxygen species, which may result in the modulation of vascular contractility⁸. Consistent with this is the notion that allopurinol, a xanthine oxidase inhibitor, not only reduces the serum UA levels but also improves vascular endothelial function in patients with chronic heart failure⁹.

Another possible explanation is that hyperuricemia may induce endothelial dysfunction by decreasing the production of nitric oxide in the vascular endothelial cells¹⁰. Adenosine synthesized locally by vascular smooth muscle in cardiac tissue is rapidly degraded by the endothelium to UA, which undergoes rapid efflux to the vascular lumen due to low intracellular pH and negative membrane potential⁷. Uric acid synthesis is increased in vivo under ischemic conditions, and therefore elevated serum UA may act as a marker of underlying tissue ischemia.

In conclusion, elevated serum UA may be a marker of local and systemic tissue ischemia and provides one possible explanation for a non-causal associative link between hyperuricemia and cardiovascular disease. Thus our study proves that SUA can be used as a marker for increased risk of acute MI. Furthermore, SUA can also be used for risk stratification after acute MI.

V. CONCLUSION:

Serum uric levels are correlated with killip's classification. Patients of higher killips class have higher uric acid levels. Combination of killip's class and serum uric acid levels after acute myocardial infarction is a good predictor of mortality. Serum uric acid levels were high in the patients who died in the seven day follow up period and the patients who died were also in higher killip's class.

REFERENCES:

- [1]. Sethi KK, ed "Preface" in Coronary Artery Disease in Indians. A Global Prospective 1998: 9 pp.
- [2]. Fang J, Alderman MH. Serum uric acid and cardiovascular mortality the NHANES I epidemiologic follow-up study, 1971-1992. National Health and Nutrition Examination I Survey. JAMA 2000; 283: 2404-2410.
- [3]. Culleton BF, Larson MG, Kannel WB, Levy D. Serum uric acid and risk for cardiovascular disease and death: the Framingham Heart Study. Ann Intern Med 1999; 131: 7-13.
- [4]. Kroll K, Bukowski TR, Schwartz LM, Knoepfler D, Bassingthwaight JB. Capillary endothelial transport of uric acid in guinea pig heart. Am J Physiol 1992, 262:H420- H431.
- [5]. Deniel I, Feig, Duk-Hee Kang et al. Uric acid and Cardiovascular Risk, The New England Journal of Medicine, October 23, 2008.
- [6]. Kojima S, Sakamoto T, Ishihara M, et al. Prognostic usefulness of serum uric acid after acute myocardial infarction (Japanese Acute Coronary Syndrome Study). Am J Cardiol 2005; 96: 489-95.
- [7]. Castelli P, Condemi AM, Brambillasca C, et al. Improvement of cardiac function by allopurinol in patients undergoing cardiac surgery. J Cardiovasc Pharmacol 1995; 25: 119-25.
- [8]. DOEHNER W, SCHOENE N, RAUCHHAUS M, et al: Effects of xanthine oxidase inhibition with allopurinol on endothelial function and peripheral blood flow in hyperuricemic patients with chronic heart failure: Results from 2 placebo-controlled studies. Circulation 105:2619-2624, 2002.
- [9]. BERRY CE, HARE JM: Xanthine oxidoreductase and cardiovascular disease: Molecular mechanisms and pathophysiological implications. J Physiol 555:589-606, 2004.
- [10]. ITOH Y, MA FH, HOSHI H, et al: Determination and bioimaging method for nitric oxide in biological specimens by diamino fluorescein fluorometry. Anal Biochem 287:203-209, 2000.