Quest Journals Journal of Medical and Dental Science Research Volume 12~ Issue 6 (June 2025) pp: 88-95 ISSN(Online): 2394-076X ISSN (Print):2394-0751 www.questjournals.org



Research Paper

Diaphragmatic Injuries on The Plateau

SD Peter*, KN Ozoilo, AH. Shitta, AF Ale, MW Isichei

Department of Surgery, Jos University Teaching Hospital, Jos, Nigeria

ABSTRACT

Background: Diaphragmatic injuries are uncommon, and its management challenging as if missed can be lethal. This study evaluate our experience in the management of diaphragmatic injury in adult patients who presented to the Accident and Emergency Unit of a Teaching Hospital in Jos, Plateau state.

Objective: To determine the pattern, management modalities and treatment outcome of traumatic gastric injury in Jos, Nigeria.

Methods: The medical records of patients admitted to our facility with diaphragmatic injury from January 2013 to December 2024 were reviewed retrospectively and analyzed.

Results: Fifty-five cases were analyzed from the period. M:F was 6.9:1, with mean age at presentation of 25.56 ± 7.16 . Commonest mechanism of injury was penetrating, and the left side was the most injured part. All the patients had laparotomy through midline approach. Stomach was also injured in 83.6% and 43.6% had infective complications. One patient developed diaphragmatic hernia. Mortality rate was 5.5%, which included the only missed injury.

Conclusion:Traumatic diaphragmatic injury commonly involve young adult males, and commonly from penetrating mechanism mainly from firearms.Left side is the side most affected, and stomach is the most other organ injured. Delayed presentation, missed injury, firearm injuries and central location is associated with higher mortality. Laparotomy remains an effective modality of treatment.

Received 05 June., 2025; Revised 14 June., 2025; Accepted 16 June., 2025 © *The author(s) 2025. Published with open access at www.questjournas.org*

I. INTRODUCTION

Diaphragmatic injuries refer to trauma to the diaphragm, which can be blunt or penetrating in nature. Injury to the diaphragm is a relatively rear but serious trauma.[1, 2, 3] The diaphragm is a c-shaped structure of muscle and fibrous tissue that separates the thoracic cavity from the abdomen. [4]As such injuries to it can affect the respiratory and cardiovascular systems, as well as septic complications that can occur from gut involvement. The clinical presentation may include difficulty in breathing, chest pain and paradoxical respiration in flail chest, or with features acute shortage of blood. A few may not have any symptom at presentation and may be missed, presenting much later with hernia and its complications. [4] Being a relatively rare injury, a high index of suspicion is therefore required to detect the occult forms, as missed injuries are associated with higher morbidity and mortality. [4]

The commonest part of the diaphragm injured is the left side [1]. Injury to the central portion is associated with higher morbidity mortality.[6] The management of diaphragmatic injuries like other serious injuries follows the Advance Trauma Life Support ATLS Protocol with resuscitation before definitive treatment. The operative treatment can be open or minimally invasive, both of which could be via the chest or abdomen. [8,9]

The aim of this study was to provide insight into the etiology, operative management and outcome, including mortality of diaphragmatic injury in Jos University Teaching Hospital.

Key words: Jos University Teaching Hospital, diaphragmatic injuries, significant trauma, laparotomy.

II. METHODOLOGY

This was a retrospective study of patients who presented with diaphragmatic injury to the Accident and Emergency unit of Jos University Teaching Hospital (JUTH), Jos, Nigeria. The medical records of the patients admitted and managed for diaphragmatic injury between January 2013 and December 2024 were retrieved from our Trauma Registry, and patients' folders, and subsequently analyzed.

Diagnosis of diaphragmatic injury was made based on the history of trauma, difficulty in breathing, chest pain, anemia, chest X-ray, focused assessment sonography for trauma (FAST), and intraoperative findings of diaphragmatic laceration.

The information extracted from the patients' medical records included age at presentation, sex, mechanism of injury, duration of injury before presentation, organs injured, complications, and outcome were transferred into Excel spread sheet and analyzed using SPSS version 23. Results are presented in tables, with frequencies and percentages, means and charts.

Ethical clearance was obtained for this study.

III. RESULTS

Fifty five (55) patients presented with traumatic diaphragmatic injury, out of a total 1078 severe thoracic and abdominal trauma patients admitted through the accident and emergency unit over the 12-year period. This indicate an average of 4.6 patients per year with traumatic diaphragmatic injury. The demographic characteristics of the patients are as shown in Tables 1 and 2, and Figure 1, with 48(87.3%) being males while 7(12.7%) were females giving an M: F of 6.9:1. Thirty patients (54.6%) were between the ages of 20-29 years and 14(25.5%) between 30-39 years accounting for the highest proportion. For both sexes the age range of 20 -29 had the highest frequency. The mean age at presentation was 25.56±7.16. Mechanism of injury was predominately penetrating as shown in Figure 2 and Tables 3a and 3b, with gunshot injury alone accounting for 31(56.4%), and stab injury 18(32.7%). Distribution of side affected were 37(67.3%) on the left, 17(30.9%) right and 1(1.8%), centrally as shown in Figure 3. All the patients had injuries to other structures, with the stomach being the most frequent abdominal organ injured having 46 patients (83.6%), followed by the small intestine with (12.7%); while the lungs was the most on the thoracic side with 9(16.4%) of the total, as shown in Table 4. Forty people, representing 72.7% of the patients presented within 9 hours while 14.5% presented after 24 hours as seen in Figure 4 and Table 5. The duration of hospital stay is as shown in Figure 4, with a mean of 19.4 days. Table 6shows and compares duration of hospital stay to mechanism of injury while Table 7 compares mechanism of injury to outcome. The commonest complication is surgical site infection as shown in Table 8, and the mortality rate is 5.5 as shown in Figure 5.

Table 1: Age distribution of patents			
Age group (vears)			Mean±SD
rige group (years)	Frequency	Percent	
< 20	9	16.4	
20-29	30	54.5	
30-39	14	25.5	
40+	2	3.6	
Total	55	100.0	25.56±7.16

Table 2: Age and Sex Distribution of patient					
Age group (years) —		Sex		x ²	Р
0010)	Male	Female	Total	<i>n</i>	
< 20	8(16.7)	1(14.3)	9(16.4)		
20-29	26(54.2)	4(57.1)	30(54.5)		
30-39	12(25.0)	2(28.6)	14(25.5)		
40+	2(4.2)	0(0.0)	2(3.6)		
Total	48(100.0)	7(100.0)	55(100.0)	0.559	0.999



Figure 1: Sex distribution of patients



Figure 2: Distribution by Etiology

Mechanism	Frequency (f)	Percent (%)
Penetrating	49	89.1
Blunt	6	10.9
Total	55	100.0

Table 3b: Distribution by etiology

Mechanism	Frequency (f)	Percent (%)
Stab	18	32.7
Gunshot	31	56.4
Blunt	6	10.9
Total	55	100.0



Figure 3: Side of lesion

Table 4a: Organs injured	
Frequency	

Organs injured	Frequency	Percent
Stomach	46	83.6
Lungs	9	16.4
Small intestine	7	12.7
Spleen	4	7.3
Transverse colon	9	16.4
Liver	7	12.7
Gall bladder	2	3.6
Kidney	1	1.8
Right femur	2	3.6
Spinal Cord	1	1.8

Table 4b: Organs injured by Mechanism

Oncons		Mec	hanism	
Organs	Penetrating	Blunt	Total	p-value
				_
Stomach	44(88.0)	6(12.0)	50(100.0)	0.999
Small intestine	7(100.0)	0(0.0)	7(100.0)	0.999
Spleen	3(75.0)	1(25.0)	4(100.0)	0.379
Transverse colon	8(88.9)	1(11.1)	9(100.0)	0.999
Liver	7(100.0)	0(0.0)	7(100.0)	0.999
Gall bladder	2(100.0)	0(0.0)	2(100.0)	0.999
Kidney	1(100.0)	0(0.0)	1(100.0)	0.999
Right femur	2(100.0)	0(0.0)	2(100.0)	0.999
Spinal Cord	1(100.0)	0(0.0)	1(100.0)	0.999



Figure 4: Duration of injury

Table 5: Duration of inj	ary vs Mechanism of injury.
--------------------------	-----------------------------

	Mechanism		
Duration	Penetrating	Blunt	Total
1-4	16(94.1)	1(5.9)	17(100.0)
5-9	21(91.3)	2(8.7)	23(100.0)
10-14	2(50.0)	2(50.0)	4(100.0)
15-19	2(100.0)	0(0.0)	2(100.0)
20-24	1(100.0)	0(0.0)	1(100.0)
25-29	1(100.0)	0(0.0)	1(100.0)
30-34	1(100.0)	0(0.0)	1(100.0)
35+	5(83.3)	1(16.7)	6(100.0)
Total	49(89.1)	6(10.9)	55(100.0)

 χ^2 (Fisher's) = 5.085, p = 0.333

Table 6: Duration of hospital stay Vs. Mechanism

		Mechanism	
Duration (Days)	Penetrating	Blunt	Total
≤7	5(83.3)	1(16.7)	6(100.0)
8-14	17(89.5)	2(10.5)	19(100.0)
15-21	10(90.9)	1(9.1)	11(100.0)
22-28	6(85.7)	1(14.3)	7(100.0)
29-35	6(100.0)	0(0.0)	6(100.0)
36+	5(83.3)	1(16.7)	6(100.0)
Total	49(89.1)	6(10.9)	55(100.0)

 χ^2 (Fisher's) = 2.037, p = 0.999

Table 7: Mechanism Vs Outcome			
Mechanism		Outcome	
	Died	Alive	Total
Penetrating	2(4.1)	47(95.9)	49(100.0)
Blunt	1(16.7)	5(83.3)	6(100.0)
Total	3(5.5)	52(94.5)	55(100.0)

 χ^2 (Fisher's) p = 0.298

Table 8: Complications			
Complications	Frequency	Percentage	
Non	25	45.5	
Surgical site infection	20	36.4	
Subphrenic abscess	1	1.8	
External gastrointestinal fistula	2	3.6	
c	1	1.8	
Burst abdomen			
Empyema	1	1.8	
Diaphragmatic hernia	1	1.8	
Paraplegia	1	1.8	
Death	3	5.5	
Total	55	100	



Figure 5: Outcome

IV. DISCUSSION

Diaphragmatic injuries are relatively uncommon but serious injuries occurring in 0.8 to 15% of all chest and abdominal trauma. [9,10] In this study we found 55 patients with diaphragmatic injuries, out of 1078 significant chest and abdominal injuries over a 12 year period, representing 5.1% of all combined severe chest and abdominal injuries. This shows an average of 4.6 patients per year which is higher than the 1.3% reported from Dakar, but lower than the 5.5% reported from Mali. [9,11] There were more males affected with M: F of 6:9. Peak age affected was between ages 20 to 29 years. Thus the age and sex distribution are in keeping with literature of severe trauma affecting young adult males. [12-15]

Left side and left dome of the diaphragm was mainly affected as shown in Figure 3. This may be because of a number of issues which include protective effect of the liver on the right side, areas of congenital weakness on the left, and the fact that left-sided injuries are easier to detect on X-rays.[1] Up to 32.7% of the etiology is from stab injuries as shown in Table 3b, and since most assailants are right handed we opine that it may also explain or contribute to the higher rate on left side. [16]

Penetrating injury was the predominant mechanism in this study, with blunt injury accounting for 10.9% as shown in Table 3b. This also is in keeping with most studies. [3,17]Gunshot injuries accounted for 56.4% of the patients, which is rather worrying in a civilian population during peace time. We opine that this may be due to increasing proliferation of light arms in circulation in our city. [14, 15] Also the high number of ethno-religious crisis in the region within the study period increased the number of gunshot injuries. [18,19] However the finding is in keeping with studies from most centers. [20,21]

There was no isolated diaphragmatic injury in our series as the diaphragm was injured along with other structures, with the stomach being the most frequent and the lungs the most on the thoracic side as shown in Table 4a. Expectedly the risk of injury to other structures was higher in penetrating injuries and even then with gunshot subset of the penetrating mechanism as indicated in Table 4b.

Duration of injury with the highest number of patients (72.7%) before presentation and intervention was between 5 to 9 hours for both blunt and penetrating injuries as shown in Figure 4. We expected the penetrating injuries to present earlier than the blunt ones as is in the literature, but there was no statistical significance in our study as seen in Table 5. [8,9,10,17]

All our patients had laparotomy for definitive treatment of the diaphragmatic injury mainly because all the patients presented with abdominal symptoms and signs, and the readily available focused assessment with sonography for trauma (FAST). In addition, most of the thoracic component of this injury could be managed by chest tube thoracostomy which all our patients had. [7-11]

Most of the complications were infective, seen in 45.4% of the patients, the commonest being wound infection in 20(36.4%) patients as shown in Table 8. This can be explained by the presence of luminal gut perforations and subsequent peritonitis in most of the patients, and the penetrating mechanisms of the injuries making these wounds at the least untidy by Rank and Wakefield classification, with a high infection rate. [22] Diaphragmatic hernia occurred in a patient which was repaired via the abdominal route. Another patient with lung parenchymal injury developed empyema thoraces, and had thoracotomy with decortication of the empyema.

Mortality rate in this study is 5.5%. This is similar to findings of Thiam O. et al in 2016[9], and lower than in most other studies. [3,7,10,12,17] Though comparing the mechanism of injury with mortality was not statistically significant (P = 0.298) as shown in Table 8, it is noteworthy to state that the only mortality from blunt injury in this study occurred in the only missed diagnosis of diaphragmatic injury in this study.

V. CONCLUSION

Traumatic diaphragmatic injury commonly involve young adult males, and commonly from penetrating mechanism mainly from firearms. Left side is the side most affected, and stomach is the most other organ injured. Delayed presentation, missed injury, firearm injuries and central location is associated with higher mortality. Laparotomy remains an effective modality of treatment.

REFERENCES

- Simon LV, Lopez RA, Burns B. Diaphragm Rupture. [Updated 2025 Apr 3]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2025 Jan-. Available from: https://www.ncbi.nlm.nih.gov/books/NBK470214/
- [2]. Corbellini C, Costa S, Canini T, Villa R, ContessiniAvesani E. Diaphragmatic rupture: A single-institution experience and literature review. Ulus TravmaAcilCerrahiDerg. 2017 Sep;23(5):421-426.
- [3]. Fair KA, Gordon NT, Barbosa RR, Rowell SE, Watters JM, Schreiber MA. Traumatic diaphragmatic injury in the American College of Surgeons National Trauma Data Bank: a new examination of a rare diagnosis. Am J Surg. 2015 May;209(5):864-8; discussion 868-9.
- [4]. Bains KNS, Kashyap S, Lappin SL. Anatomy, Thorax: Diaphragm. [Updated 2023 Jul 24]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2025 Jan-. Available from: https://www.ncbi.nlm.nih.gov/books/NBK519558/
- [5]. Kumar A, Bagaria D, Ratan A, Gupta A. Missed diaphragmatic injury after blunt trauma presenting with colonic strangulation: a rare scenario. BMJ Case Rep. 2017 Aug 07;2017.
- [6]. Nwafor IA, Eze JC, Aminu MB. Traumatic diaphragmatic rupture through the central tendon with herniation of the stomach and coils of small bowel into the pericardial cavity. Niger J Med. 2011 Oct-Dec; 20(4):492-3.
- [7]. Lopez PP, Arango J, Gallup TM, Cohn SM, Myers J, Corneille M, et al. Diaphragmatic injuries: what has changed over a 20-year period? Am Surg 2010; 76:512–6.
- [8]. Kim Y. A clinical analysis of traumatic diaphragmatic injury. J Korean SocTraumatol 2004; 17:67–73.
- [9]. Thiam, O., Konate, I., Gueye, M.L. et al. Traumatic diaphragmatic injuries: epidemiological, diagnostic and therapeutic aspects. SpringerPlus, 5, 1614 (2016).
- [10]. Mihos P, Potaris K, Gakidis J, Paraskevopoulos J, Varvatsoulis P, Gougoutas B et al (2003) Traumatic rupture of the diaphragm: experience with 65 patients. InjInt J Care Inj 34:169–172
- [11]. Traoré, A., Konaté, M., Diarra, A., Tounkara, I., Traoré, M., Doumbia, A., Bah, A., Sidibé, B., Maïga, A., Koné, T., Karembé, B., Saye, Z., Kéita, K., Bouaré, Y., Koné, A., Diakité, I., Kanté, L., Dembélé, B., Traoré, A. and Togo, A. (2022) Traumatic Diaphragmatic Injury at Gabriel Toure University Hospital, Mali. Surgical Science, 13, 110-118.
- [12]. Søreide K. Epidemiology of major trauma. Journal of British Surgery. 2009;96(7):697-8.
- [13]. Alberdi F, Garcia I, Atutxa L, Zabarte M. Epidemiology of severe trauma. MedicinaIntensiva (English Edition). 2014;38(9):580-8.
- [14]. Sakran JV, Greer SE, Werlin E, McCunn M. Care of the injured worldwide: trauma still the neglected disease of modern society. Scandinavian journal of trauma, resuscitation and emergency medicine. 2012; 20(1):1-6.

- [15]. Păun S, Beuran M, Negoi I, Runcanu A, Gaspar B. Trauma-epidemiology: where are we today?]. Chirurgia (Bucur). 2011 Jul-Aug; 106(4):439-43.
- [16]. Eze, Joy N., E. N. Obikili, Udechukwu F Ezepue, O F N Ozoemena and Emmanuel A. Esom. "Patterns of Handedness and Sociocultural Influences on Dextrality Amongst University of Nigeria Medical Students." (2010).
- [17]. Lee, Sang Su et al. "Comparison of Penetrating and Blunt Traumatic Diaphragmatic Injuries." *Journal of Trauma and Injury* vol. 32,4 (2019): 210-219
- [18]. Osaretin I, Akov E. Ethno-religious conflict and peace building in Nigeria: The case of Jos, Plateau State. Academic Journal of Interdisciplinary Studies. 2013; 2(1):349-352.
- [19]. John IA, Mohammed AZ, Pinto AD, Nkanta CA. Gun violence in Nigeria: A focus on ethno-religious conflict in Kano. Journal of public health policy. 2007;28(4):420-31
- [20]. Inci I, Ozçelik C, Taçyildiz İ, Nizam O, Eren N, Ozgen G. Penetrating chest injuries: unusually high incidence of high-velocity gunshot wounds in civilian practice. World J Surg. 1998 May. 22 (5):438-42.
- [21]. Onat S, Ulku R, Avci A, Ates G, Ozcelik C. Urgent thoracotomy for penetrating chest trauma: analysis of 158 patients of a single center. *Injury*. 2011 Sep. 42 (9):900-4.
- [22]. Mohil R. Classification of wounds. Principles and practice of wound care. 2012; 1:42-52.