



Research Paper

Prevalence of premature greying of hair and Its Association with Lifestyle and Dietary Factors Among Young Adults: A Cross-Sectional Study

Dr. Yogita Kotwani¹, Dr. Prashant Solanke², Dr. Rajesh Saraf³

¹ Intern, Department of Research, Dr. Ulhas Patil Medical College, Jalgaon, Maharashtra, India, 425309.
| Phone: 8390034594

² Dean and Professor of Community Medicine, Department of Community Medicine, Dr. Ulhas Patil Medical College, Jalgaon, Maharashtra, India, 425309.
Phone: 97519 26130

³ Assistant Professor, Department of Paediatrics, Dr. Ulhas Patil Medical College, Jalgaon, Maharashtra, India.
Phone: 9422182309

Corresponding Author: Dr. Yogita Kotwani, Intern, Department of Research, Dr. Ulhas Patil Medical College, Jalgaon, Maharashtra, India - 425309.

ABSTRACT:

Background: Premature greying of hair is a common hair-pigmentation disorder that may occur at a relatively young age. Its development is considered multifactorial, with genetic, environmental, nutritional and lifestyle-related factors being implicated.

Objectives:

1. To determine the prevalence of premature greying of hair among young adults.
2. To assess the association between lifestyle factors and premature greying of hair among young adults.
3. To assess the association between dietary factors and premature greying of hair among young adults.

Methodology: A cross-sectional study was conducted using a structured questionnaire. The analysis was restricted to participants aged 18–30 years. The sample size was calculated using $n = Z^2pq/d^2$, taking $Z = 1.96$, $p = 27.3\%$, $q = 72.7\%$ and an absolute allowable error of 10%, giving a minimum sample size of approximately 77 participants. Of the available responses, 87 participants aged 18–30 years were included in the final analysis. Premature greying was operationally defined as currently having grey/white hair with onset before 25 years of age. Associations were assessed using Pearson's chi-square test or Fisher's exact test, as appropriate. A p -value < 0.05 was considered statistically significant.

Results: Among the 87 participants aged 18–30 years, 62 (71.3%) reported currently having grey/white hair. Using the predefined operational definition, 56 participants (64.4%) were classified as having premature greying. Sleep duration was significantly associated with premature greying ($p = 0.044$), as was frequency of perceived stress ($p = 0.026$). Among dietary factors, frequency of fruit consumption was significantly associated with premature greying ($p = 0.049$). Other assessed lifestyle and dietary factors were not statistically significant.

Conclusion: Premature greying was frequently reported in the study sample. Sleep duration, perceived stress and fruit consumption showed statistically significant associations with premature greying, while the other assessed factors did not.

KEYWORDS: Premature greying; lifestyle; dietary factors; young adults.

Received 07 Oct., 2026; Revised 16 Oct., 2026; Accepted 18 Oct., 2026 © The author(s) 2026.
Published with open access at www.questjournals.org

I. INTRODUCTION

Hair graying is a visible manifestation of the aging process and occurs because of a reduction or cessation of melanin production by melanocytes within the hair follicle. Although graying is generally

associated with increasing age, premature graying may occur much earlier in life.¹

Premature graying of hair, also referred to as premature canities, has been described using age-based definitions that vary according to ethnicity. In Asian populations, graying before approximately 25 years of age is commonly considered premature.^{2,3}

The development of premature graying is considered multifactorial. Genetic predisposition and family history are important factors, while oxidative stress, smoking, nutritional deficiencies and certain medical conditions have also been reported in the literature.^{1,2} A systematic review and meta-analysis reported associations between premature graying and smoking, deficiencies of vitamin B12, folic acid and biotin, low serum calcium and ferritin, family history, lack of exercise, thyroid disorders and other metabolic conditions.²

Lifestyle factors may also influence hair health. Smoking, for example, has been investigated in relation to premature hair graying, with proposed mechanisms involving oxidative stress and vascular effects.⁴ Psychological stress has also been proposed as a possible contributor to hair pigmentation changes, although the relationship is complex and requires further investigation.

Diet and nutritional status are also of interest because several micronutrients are involved in melanogenesis and normal hair follicle function. Deficiencies involving vitamin B12, folate, iron, calcium, copper and other micronutrients have been reported in association with premature graying.^{1,2}

An Indian community-based cross-sectional study among 384 young adults aged 15–30 years reported a prevalence of premature graying of 27.3%. The study also reported significant associations with paternal and maternal history, smoking, sunlight exposure and body mass index.⁵

Despite the increasing observation of premature graying among younger individuals, there remains a need to examine potentially modifiable lifestyle and dietary factors in different populations. Therefore, the present study was undertaken to determine the prevalence of premature graying and assess its association with selected lifestyle and dietary factors among adults aged 18–30 years.

II. OBJECTIVES

1. To determine the prevalence of premature greying of hair among young adults.
2. To assess the association between lifestyle factors and premature greying of hair among young adults.
3. To assess the association between dietary factors and premature greying of hair among young adults.

III. METHODOLOGY

Study design

A cross-sectional study was conducted to assess the prevalence of premature greying of hair and its association with selected lifestyle and dietary factors.

Study setting

The study was conducted among the young adults aged 18 to 30 years from Vidarbha and Khandesh region, Maharashtra, India.

Study period

1 August 2026 to 1 October 2026. Study population

The study population comprised individuals who participated in the structured questionnaire-based data collection. The final analysis was restricted to participants aged 18–30 years.

Sample size

The required sample size was calculated using $n = Z^2pq/d^2$, where $Z = 1.96$ at a 95% confidence level, $p = 27.3\% = 0.273$, $q = 100p = 72.7\% = 0.727$, and $d = 10\% = 0.10$. The calculated minimum sample size was 76.24, approximately 77 participants. A total of 87 participants aged 18–30 years were available and included in

The prevalence value used for sample-size planning was 27.3%, based on the prevalence reported in a comparable community-based Indian study among young adults.⁵

Sampling technique

A community-based non-random convenience sampling technique was used for participant recruitment through the structured questionnaire.

Inclusion criteria

1. Participants aged 18–30 years.
2. Participants who provided responses to the study questionnaire.

3. Participants who provided information regarding their hair-graying status and relevant lifestyle and dietary factors.

Exclusion criteria

Participants with incomplete responses that prevented assessment of premature graying status were excluded from the analysis. Critically ill participants or those unable to provide reliable questionnaire responses were excluded.

Data collection

Information was collected using a structured questionnaire covering sociodemographic characteristics, hair graying, lifestyle practices and dietary habits.

Premature greying of hair was operationally defined as self-reported current grey/white hair with first noticing of grey/white hair before 25 years of age. This age threshold is consistent with commonly used definitions for Asian populations.³

Statistical analysis

Categorical variables were summarized as frequencies and percentages. Associations between premature greying and categorical lifestyle or dietary factors were assessed using Pearson's chi-square test or Fisher's

Ethical considerations

Participation was based on voluntary completion of the study questionnaire. Participant responses were treated confidentially and used for research purposes. Institutional Ethics Committee approval was obtained prior to commencement of the study.

IV. RESULTS

Study participants: A total of 87 participants aged 18–30 years were included in the final analysis. The mean age was 24.0 ± 2.7 years, with a median age of 24 years and an age range of 18–30 years. There were 49 females (56.3%) and 38 males (43.7%). Most participants were from urban areas (51; 58.6%), followed by semi-urban (22; 25.3%) and rural (14; 16.1%). Students constituted 64 participants (73.6%), followed by

Table 1. Sociodemographic characteristics of study participants (N = 87)

1. Most participants were aged 18–24 years (62.1%) and female (56.3%).
2. The majority were urban residents (58.6%) and students (73.6%).

Characteristic	Category	n	%
Age group	18–24 years	54	62.1
	25–30 years	33	37.9
Sex	Female	49	56.3
	Male	38	43.7
Residence	Urban	51	58.6
	Semi-urban	22	25.3
	Rural	14	16.1
Occupation	Student	64	73.6
	Employed	18	20.7
	Other	5	5.7

Prevalence of premature greying

Of the 87 participants, 62 (71.3%) reported current grey/white hair, 16 (18.4%) reported no grey/white hair and 9 (10.3%) were unsure. Using the predefined operational definition, 56 participants (64.4%) were classified as having premature greying and 31 (35.6%) were not classified as having premature greying.


Table 2. Prevalence of current grey/white hair and premature greying (N = 87)

1. Current grey/white hair was reported by 62 participants (71.3%).
2. Premature greying was classified in 56 participants (64.4%).

Measure	Category	n	%
Current grey/white hair	Yes	62	71.3
	No	16	18.4
	Not sure	9	10.3
Premature greying	Yes	56	64.4
	No/not classified	31	35.6

Figure 1. Premature greying among participants aged 18-30 years

Prevalence of premature greying among participants aged 18-30 years



Association between lifestyle factors and premature greying

Using Pearson's chi-square test or Fisher's exact test, as appropriate, sleep duration was significantly associated with premature greying ($p = 0.044$), as was frequency of perceived stress ($p = 0.026$). Physical activity, tobacco use, alcohol use, heat-styling tools, chemical hair treatment and water hardness were not statistically significant ($p > 0.05$).

Table 3. Association between lifestyle factors and premature greying of hair

1. Sleep duration ($p = 0.044$) and perceived stress ($p = 0.026$) showed statistically significant associations.
2. Other assessed lifestyle factors were not statistically significant ($p > 0.05$).

Lifestyle factor	Category	Premature greying n/N (%)	p-value
Sleep duration	<6 h	20/38 (52.6%)	0.044
	≥6 h	36/49 (73.5%)	
Perceived stress	Never/rarely	7/17 (41.2%)	0.026
	Moderate/high	49/70 (70.0%)	
Physical activity	<3 days/week	48/73 (65.8%)	0.538
	≥3 days/week	8/14 (57.1%)	
Tobacco use	Any use	4/6 (66.7%)	1.000
	No use	52/81 (64.2%)	
Alcohol use	No	49/77 (63.6%)	1.000
	Yes	7/10 (70.0%)	
Heat-styling tools	No	29/47 (61.7%)	0.574
	Yes	27/40 (67.5%)	
Chemical hair treatment	No	47/69 (68.1%)	0.153
	Yes	9/18 (50.0%)	
Water hardness	Don't know	6/11 (54.5%)	0.176
	Hard	40/56 (71.4%)	
	Soft	10/20 (50.0%)	

Association between dietary factors and premature greying

Frequency of fruit consumption showed a statistically significant association with premature greying ($p = 0.049$). The other dietary factors assessed were not statistically significant ($p > 0.05$).

Table 4. Association between dietary factors and premature greying of hair

1. Fruit consumption was significantly associated with premature greying ($p=0.049$).
2. The other assessed dietary factors were not statistically significant ($p>0.05$).

Dietary factor	Category	Premature greying n/N(%)	p-value
Meals/day	1-2	40/63(63.5%)	0.782
	≥ 3	16/24(66.7%)	
Green leafy vegetables	<3/week	48/72(66.7%)	0.327
	≥ 3 /week	8/15(53.3%)	
Fruits	<3/week	52/76(68.4%)	0.049
	≥ 3 /week	4/11(36.4%)	
Pulses/beans/lentils	<3/week	48/71(67.6%)	0.184
	≥ 3 /week	8/16(50.0%)	
Milk/dairy products	<3/week	43/64(67.2%)	0.360
	≥ 3 /week	13/23(56.5%)	
Animal-protein foods	<3/week	49/75(65.3%)	0.638
	≥ 3 /week	7/12(58.3%)	
Fast food/junk food	≤ 2 /week	38/54(70.4%)	0.135
	≥ 3 /week	18/33(54.5%)	
Skipping meals	Never/rarely	23/34(67.6%)	0.609
	Sometimes/often	33/53(62.3%)	
Special diet	No	26/44(59.1%)	0.299
	Yes	30/43(69.8%)	
Vitamin/minerals supplements	No	47/74(63.5%)	0.764
	Yes	9/13(69.2%)	

Table 4. Association between dietary factors and premature greying of hair (continued)

Dietary factor	Category	Premature greying n/N(%)	p-value
Nutritional deficiency	Don't know	5/7(71.4%)	0.890
	No	41/65(63.1%)	
	Yes	10/15(66.7%)	

Table 5. Statistically significant associations with premature greying

Factor	Comparison	Premature greying n/N(%)	p-value
Sleep duration	<6h vs ≥ 6 h	20/38(52.6%) vs 36/49(73.5%)	0.044
Perceived stress	Never/rarely vs Moderate/high	7/17(41.2%) vs 49/70(70.0%)	0.026
Fruit consumption	<3/week vs ≥ 3 /week	52/76(68.4%) vs 4/11(36.4%)	0.049

DISCUSSION

Premature graying of hair is considered a multifactorial phenomenon involving genetic, biological, nutritional and environmental influences.^{1,2} The operational definition used in this study is consistent with commonly used definitions for Asian populations.³ Lifestyle factors, including smoking, have also been investigated in relation to premature hair graying.

The observed prevalence in the present dataset is higher than the 27.3% reported in a community-based study conducted among 384 young adults aged 15–30 years in Srinagar, Uttarakhand.⁵ Differences may relate to participant characteristics, sampling methods, outcome definitions and methods of assessment. The earlier study assessed graying through direct scalp examination, whereas the present study used self-reported questionnaire data.⁵

Sleep duration and perceived stress were significantly associated with premature greying in the present study. These findings indicate associations within this sample and do not establish that inadequate sleep or stress causes premature greying.

Previous research has reported associations between premature graying and several lifestyle and medical factors, including smoking and lack of exercise.⁴ The present study did not demonstrate statistically significant associations with physical activity, tobacco use or alcohol consumption.

Frequency of fruit consumption was significantly associated with premature greying, whereas the other dietary factors assessed were not statistically significant. This finding should be interpreted cautiously because the study relied on self-reported dietary frequency and did not include laboratory measurements of nutritional status.

Family history and genetic susceptibility are also important considerations. Previous Indian research has reported associations between family history and premature greying.³ The present analysis did not use family history as one of the primary lifestyle/dietary association tables, and therefore no independent conclusion regarding family history is drawn here.

CONCLUSION

Premature greying was identified in 56 of 87 participants (64.4%) using the study's operational definition.

Current grey/white hair was reported by 62 participants (71.3%).

Sleep duration showed a statistically significant association with premature greying ($p = 0.044$).

Perceived stress was also significantly associated with premature greying ($p = 0.026$).

Frequency of fruit consumption showed a statistically significant association with premature greying ($p = 0.049$).

The other assessed lifestyle factors, including physical activity, tobacco use, alcohol use, heat-styling tools, chemical hair treatment and water hardness, were not statistically significant.

The other assessed dietary factors were also not statistically significant.

The findings highlight sleep duration, perceived stress and fruit-consumption frequency as factors associated with premature greying in this study sample.

These findings suggest that sleep duration, perceived stress and fruit-consumption frequency may be relevant factors associated with premature greying in this study population.

LIMITATIONS

1. The study used a cross-sectional design; therefore, causal relationships cannot be established.
2. The sample size was relatively small, although the final sample of 87 exceeded the calculated minimum of 77.
3. Hair greying and lifestyle/dietary factors were assessed using self-reported questionnaire responses.
4. The study did not include laboratory assessment of nutritional biomarkers, and the findings may not be generalizable beyond the study population.

FUNDING

No external funding was received for this study.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

ACKNOWLEDGEMENT

The authors would like to thank all the participants who took part in the study and provided the required information for data collection.

REFERENCES

1. Kumar AB, Shamim H, Nagaraju U. Premature greying of hair: review with updates. *Int J Trichol*. 2018;10(5):198-203.
2. Mahendiratta S, Sarma P, Kaur H, et al. Premature greying of hair: risk factors, co-morbid conditions, pharmacotherapy and reversal? a systematic review and meta-analysis. *Dermatol Ther*. 2020;33(6):e13990. doi:10.1111/dth.13990.
3. Pandhi D, Khanna D. Premature greying of hair. *Indian J Dermatol Venereol Leprol*. 2013;79:641-653.
4. Trüeb RM. Aging of hair. *J Cosmet Dermatol*. 2005;4(2):60-72. doi:10.1111/j.1473-2165.2005.40203.x.
5. Nath B, Gupta V, Kumari R. A community based study to estimate prevalence and determine correlates of premature greying of hair among young adults in Srinagar, Uttarakhand, India. *Int J Trichol*. 2020;12(5):206-212. doi:10.4103/ijt.ijt_28_19.
6. Chakrabarty S, Krishnappa PG, Gowda DG, Hiremath J. Factors associated with premature hair greying in a young Indian population. *Int J Trichol*. 2016;8(1):11-14.
7. Zayed AA, Shahait AD, Ayoub MN, Yousef AM. Smokers' hair: does smoking cause premature hair greying? *Indian Dermatol Online J*. 2013;4(2):90-92.
8. Trüeb RM. Oxidative stress in aging of hair. *Int J Trichology*. 2009;1(1):6-14. doi:10.4103/0974-7753.51923.
9. Poonia K, Bhalla M. Premature Graying of Hair: A Comprehensive Review and Recent Insights. *Indian Dermatol Online J*. 2024;15(5):721-731. doi:10.4103/idoj.idoj_807_23.
10. Desai DD, Karim M, Nohria A, et al. Premature hair greying: a multifaceted phenomenon. *Int J Dermatol*. 2025;64(5):819-829.