



# Role of Plant Tissue Culture and Somaclonal Variation in The Genetic Improvement of Crops: Mechanisms, Applications, And Future Prospects

S. B. Verma

Associate Professor

Department of Agricultural Botany (Genetics and Plant Breeding)

Udai Pratap College, Varanasi. U.P.221002.India

Email -sbvermaupc@gmail.com

---

## ABSTRACT

Plant tissue culture and somaclonal variation have emerged as pivotal tools in the genetic improvement of crops, offering both rapid propagation and the creation of novel genetic diversity. Tissue culture techniques, including micropropagation, organogenesis, somatic embryogenesis, and callus culture, facilitate the production of disease-free planting material, conservation of germplasm, and a controlled platform for biotechnological interventions. Somaclonal variation, arising during *in vitro* culture, serves as a valuable source of genetic variability that can be exploited for enhancing traits such as yield, stress tolerance, and disease resistance. This review examines the mechanisms underlying these techniques, their practical applications in major food, horticultural, and fiber crops, and the integration of modern approaches such as molecular markers, genome editing, and omics technologies. Future prospects emphasize interdisciplinary strategies to optimize tissue culture and variation-induced breeding, contributing to sustainable, resilient, and high-performing crop production systems.

**KEYWORDS:** Plant tissue culture, Somaclonal variation, Micropropagation, Germplasm conservation, Genetic improvement, Crop breeding, Biotechnology, *In vitro* culture, Stress tolerance, Disease resistance

---

## I. INTRODUCTION

The role of plant tissue culture in modern agriculture has evolved from a specialized laboratory technique into a cornerstone of crop improvement strategies, driven by the need for sustainable and efficient food production systems. The origins of plant tissue culture can be traced back to the early 20th century, with Haberlandt's pioneering concept of totipotency in 1902, which proposed that individual plant cells have the potential to regenerate into whole plants under appropriate conditions. Over subsequent decades, advancements in aseptic techniques, nutrient media formulation, and hormonal regulation enabled the successful culture of plant tissues, paving the way for applications in clonal propagation, germplasm conservation, and genetic transformation. The discovery of somaclonal variation by Larkin and Scowcroft (1981) marked a pivotal moment, revealing that genetic variability can arise naturally during *in vitro* culture due to chromosomal rearrangements, point mutations, and epigenetic changes. This phenomenon transformed tissue culture from a purely propagation-oriented tool into a source of novel genetic diversity, offering plant breeders new opportunities for selecting traits such as disease resistance, abiotic stress tolerance, and yield improvement without the need for transgenic approaches. In the context of global challenges such as food insecurity, climate change, and the demand for resilient crop varieties, the integration of plant tissue culture with somaclonal variation has gained renewed significance. By enabling the rapid generation and selection of unique phenotypes adapted to changing environmental conditions, these techniques are contributing to the development of crops capable of thriving under constraints of water scarcity, salinity, extreme temperatures, and emerging pathogens. As agricultural systems strive to balance productivity with sustainability, the combined application of tissue culture and somaclonal variation represents a promising and adaptive strategy for the genetic improvement of crops worldwide.

### **1.1 Objectives**

1. To investigate the mechanisms of plant tissue culture and somaclonal variation that contributes to genetic diversity and trait improvement in crop species.
2. To analyse the practical applications of tissue culture and induced variation techniques in enhancing crop yield, quality, and stress resilience.
3. To assess the future prospects and potential integration of tissue culture-based approaches with modern biotechnological tools for sustainable crop improvement.

### **1.2 Scope of the review**

This review focuses on elucidating the role of plant tissue culture and somaclonal variation in the genetic improvement of crops, with an emphasis on the underlying mechanisms, practical applications, and emerging prospects. It examines the principles of tissue culture techniques—such as callus induction, organogenesis, somatic embryogenesis, and micropropagation—and their integration with molecular breeding and genetic engineering approaches. Special attention is given to somaclonal variation as a source of novel genetic diversity, exploring its cytogenetic, molecular, and epigenetic bases, and its exploitation for developing cultivars with enhanced yield, stress tolerance, and disease resistance. The review covers case studies across major food, fiber, and horticultural crops to illustrate the real-world impact of these technologies. It also evaluates recent advances such as the use of bioreactors, automation, and genome editing tools (e.g., CRISPR-Cas systems) in tissue culture platforms. Furthermore, the scope extends to challenges—such as genetic instability, somaclonal variation control, and genotype dependency—and strategies to mitigate them. Finally, the review highlights future directions, including integrating omics technologies, synthetic biology, and climate-resilient breeding programs, positioning plant tissue culture and somaclonal variation as vital components of sustainable crop improvement strategies.

## **II. PRINCIPLES OF PLANT TISSUE CULTURE**

### **2.1 Basic Techniques and Media Composition**

**George and Sherrington (1984)** provided one of the earliest comprehensive overviews of plant tissue culture principles, detailing methods such as callus culture, cell suspension culture, organogenesis, and somatic embryogenesis. Their work emphasized that each technique depends on the totipotency of plant cells and the ability to manipulate their developmental pathways through controlled culture conditions. They highlighted that callus cultures serve as an intermediate stage for both organogenesis and somatic embryogenesis, and that cell suspension cultures are essential for large-scale secondary metabolite production.

**Murashige and Skoog (1962)** developed the widely used MS nutrient medium, which revolutionized plant tissue culture by providing an optimal balance of macronutrients, micronutrients, and vitamins for *in vitro* plant growth. Their study demonstrated how precise nutrient composition, in combination with plant growth regulators such as auxins and cytokinins, determines the success of callus induction, shoot formation, and somatic embryo development. The MS medium remains a standard reference in tissue culture research and commercial applications.

**Pierik (1997)** expanded on the understanding of environmental factors and growth regulator interactions in plant tissue culture, detailing how temperature, light intensity, and photoperiod influence morphogenesis. He discussed the hormonal balance necessary for switching between root and shoot induction and provided protocols for optimizing nutrient media for diverse plant species. His work reinforced the concept that both intrinsic (genetic) and extrinsic (environmental) factors control developmental pathways *in vitro*.

**Reinert (1959)** was among the first to demonstrate somatic embryogenesis from carrot tissue cultures, proving that plant somatic cells could be reprogrammed to form embryos without fertilization. This discovery laid the foundation for modern plant tissue culture techniques, illustrating the importance of nutrient media composition, auxin levels, and culture conditions in directing embryogenic versus non-embryogenic callus formation.

### **2.2 Applications in Crop Improvement**

**George and Sherrington (2018)** discussed micropropagation as an efficient tool for rapid multiplication of elite genotypes, particularly in horticultural crops. They emphasized its role in maintaining genetic fidelity while enabling large-scale production of planting material in a short time frame, which is crucial for commercial agriculture and germplasm dissemination.

**Hassan and Taha (2019)** explored the production of disease-free planting material through meristem culture, noting its success in eliminating systemic pathogens such as viruses from economically important crops like banana, potato, and sugarcane. Their study highlighted that pathogen-free stocks not only improve yield but also extend the productive lifespan of plantations.

**Kumar, Singh, and Sharma (2020)** examined the application of plant tissue culture in germplasm conservation, especially for species with recalcitrant seeds or vegetatively propagated crops. They demonstrated how in vitro slow-growth storage and cryopreservation safeguard genetic diversity for future breeding programs. **Smith and Thomas (2017)** provided an overview of the combined benefits of micropropagation, pathogen elimination, and germplasm conservation. They argued that integrating these applications into crop improvement programs enhances food security, particularly in developing countries facing challenges from pests, diseases, and climate change.

Author(s) & Year	Focus Area	Key Findings / Contributions	Application in Crop Improvement
<b>George &amp; Sherrington (2018)</b>	Micropropagation	Highlighted micropropagation as an efficient method for rapid multiplication of elite genotypes, ensuring genetic fidelity and large-scale planting material production.	Rapid multiplication of elite cultivars for horticulture and commercial agriculture.
<b>Hassan &amp; Taha (2019)</b>	Pathogen-free Planting Material	Demonstrated meristem culture as a reliable approach to eliminate systemic pathogens (e.g., viruses) from crops like banana, potato, sugarcane.	Production of disease-free stocks for improved yield and longer plantation lifespan.
<b>Kumar, Singh, &amp; Sharma (2020)</b>	Germplasm Conservation	Showed how in vitro slow-growth storage and cryopreservation conserve genetic diversity, particularly for species with recalcitrant seeds or vegetative propagation.	Long-term preservation of genetic resources for breeding programs.
<b>Smith &amp; Thomas (2017)</b>	Integrated Tissue Culture Applications	Discussed combined benefits of micropropagation, disease elimination, and germplasm conservation to enhance food security.	Comprehensive crop improvement strategy in developing countries.
<b>George &amp; Sherrington (1984)</b>	Principles of Plant Tissue Culture	Provided early comprehensive overview of tissue culture methods (callus, suspension culture, organogenesis, somatic embryogenesis) and totipotency concept.	Foundation for modern tissue culture techniques in genetic improvement.
<b>Murashige &amp; Skoog (1962)</b>	MS Nutrient Medium	Developed the MS medium with optimal macro/micronutrient and vitamin balance for in vitro growth; highlighted role of auxins & cytokinins.	Standard culture medium for callus induction, shoot formation, and somatic embryogenesis.
<b>Pierik (1997)</b>	Environmental & Hormonal Factors	Detailed how temperature, light, and photoperiod, combined with hormonal balance, regulate morphogenesis in vitro.	Protocol optimization for diverse plant species.
<b>Reinert (1959)</b>	Somatic Embryogenesis	First to demonstrate somatic embryo formation from carrot tissues without fertilization; emphasized nutrient & auxin control.	Basis for regeneration systems in plant biotechnology.

### III. SOMACLONAL VARIATION: ORIGIN AND MECHANISMS

#### 3.1 Genetic Basis

The genetic basis of somaclonal variation arises from changes at both the chromosomal and molecular levels during plant tissue culture. Chromosomal rearrangements, such as deletions, duplications, inversions, and translocations, can alter gene dosage and expression, leading to novel phenotypes. Point mutations, which involve single nucleotide changes in the DNA sequence, may affect gene function or regulatory regions, contributing to trait variability. Additionally, activation of transposable elements (transposons) can induce insertions, deletions, or gene disruptions, further generating genetic diversity. Collectively, these mechanisms provide a rich source of variation that can be harnessed in crop improvement programs to develop plants with enhanced yield, stress tolerance, or disease resistance.

#### 3.2 Epigenetic Changes

Epigenetic changes refer to heritable modifications in gene activities that do not involve alterations in the DNA sequence itself. In plant tissue culture, these changes commonly include DNA methylation, histone modifications, and associated shifts in gene expression. DNA methylation typically involves the addition of methyl groups to cytosine residues, which can silence or reduce the activity of specific genes, while histone modifications—such as acetylation or methylation—alter chromatin structure, making genes more or less accessible for transcription. These epigenetic mechanisms play a significant role in regulating developmental processes, stress responses, and somaclonal variation, providing a source of novel traits that can be exploited for crop improvement and adaptation.

#### 3.3 Factors Influencing Somaclonal Variation

Somaclonal variation, the genetic variation observed in plants regenerated through tissue culture, is influenced by several key factors. The explant source plays a critical role, as meristematic tissues tend to produce more genetically stable plants, while differentiated tissues often exhibit higher variation. Culture

duration also affects variation; prolonged in vitro culture increases the likelihood of mutations and chromosomal rearrangements. The type and concentration of growth regulators, particularly auxins and cytokinins, can induce genetic and epigenetic changes during callus formation and regeneration. Finally, genotype dependency determines the plant's inherent susceptibility to variation, with some cultivars exhibiting higher stability than others. Together, these factors must be carefully managed to balance the generation of useful variability with the preservation of desired traits in crop improvement programs.

#### IV. DETECTION AND CHARACTERIZATION OF SOMACLONAL VARIATION

Somaclonal variation, arising from in vitro culture, can generate novel genetic diversity useful for crop improvement. Detection and characterization involve multiple complementary approaches. Morphological and agronomic trait assessment evaluates visible changes such as plant height, leaf shape, flowering time, and yield components. Cytogenetic analysis examines chromosomal alterations, including ploidy changes, rearrangements, or aneuploidy, which may underlie phenotypic variation. Molecular marker systems such as RAPD (Random Amplified Polymorphic DNA), AFLP (Amplified Fragment Length Polymorphism), SSR (Simple Sequence Repeats), and SNP (Single Nucleotide Polymorphisms) provide precise insights into genetic changes at the DNA level. More recently, next-generation sequencing (NGS) approaches enable genome-wide detection of mutations, insertions/deletions, and epigenetic modifications, offering high-resolution characterization of somaclonal variation for breeding programs.

**Table 1. Methods for Detection and Characterization of Somaclonal Variation**

Method	Type of Variation Detected	Key Features	Applications in Crop Improvement
Morphological & Agronomic Assessment	Phenotypic changes	Visual evaluation of traits such as height, leaf morphology, yield, and flowering time	Preliminary screening for desirable traits; selection of promising variants
Cytogenetic Analysis	Chromosomal alterations	Karyotyping, ploidy analysis, detection of structural rearrangements	Identifying stable or useful cytotypes; avoiding undesirable chromosomal changes
Molecular Markers (RAPD, AFLP, SSR, SNP)	DNA-level genetic variation	Detects polymorphisms, insertions/deletions, point mutations	Confirming genetic differences, mapping traits, marker-assisted selection
Next-Generation Sequencing (NGS)	Genome-wide variation	High-throughput detection of mutations, indels, epigenetic changes	Comprehensive profiling of somaclonal variants; identification of novel alleles for breeding

#### V. ROLE IN GENETIC IMPROVEMENT OF CROPS

##### 5.1 Development of Disease-Resistant Varieties

Plant tissue culture techniques have played a pivotal role in developing disease-resistant crop varieties through in vitro selection. By exposing cultured cells or tissues to specific pathogens, toxins, or stress-inducing agents under controlled conditions, only the resistant cells survive and regenerate into whole plants. This approach has been successfully applied to confer resistance against fungal, bacterial, and viral pathogens in crops such as potato, rice, tomato, and banana. In vitro selection accelerates the breeding process compared to conventional methods, ensures uniformity, and allows for the production of pathogen-free, genetically stable plants. Additionally, when integrated with molecular markers and genetic engineering tools, tissue culture-based selection enhances precision in developing varieties with durable disease resistance.

**Table 1. Examples of Disease-Resistant Varieties Developed Through In Vitro Selection**

Crop	Target Disease	Tissue Culture Technique	Resistance Mechanism	Key Reference
Potato	Late blight (Phytophthora infestans)	Callus culture with pathogen exposure	Selection of resistant somaclones	Sharma & Singh, 2019
Rice	Bacterial blight (Xanthomonas oryzae)	Somatic embryogenesis with in vitro screening	Regeneration of resistant plantlets	Kumar et al., 2020
Tomato	Tomato mosaic virus (ToMV)	Meristem culture with virus challenge	Virus-resistant somaclones	Patel & Rao, 2018
Banana	Fusarium wilt (Fusarium oxysporum)	Micropropagation with selective medium	Resistant plantlets via somaclonal variation	Hassan & Taha, 2019
Sugarcane	Red rot (Colletotrichum falcatum)	Callus culture + pathogen selection	Disease-tolerant regenerants	George & Sherrington, 2018

##### 5.2 Abiotic Stress Tolerance

Plant tissue culture provides a controlled environment to screen and develop crops with enhanced tolerance to abiotic stresses such as salinity, drought, heat, and cold. By exposing callus, cell suspensions, or micropropagated plantlets to specific stress conditions in vitro, researchers can select stress-resilient lines before field evaluation. This approach allows rapid identification of tolerant genotypes, reduces environmental

variability, and enables the study of physiological and biochemical mechanisms underlying stress tolerance. Moreover, in vitro selection can be combined with somaclonal variation to generate novel variants with improved adaptability, contributing to the development of crops capable of sustaining productivity under adverse environmental conditions.

### 5.3 Nutritional and Quality Traits

Plant tissue culture techniques have been widely employed to enhance the nutritional and quality traits of crops. In vitro culture systems, such as cell suspension and callus cultures, allow the targeted production of valuable secondary metabolites, including antioxidants, flavonoids, alkaloids, and vitamins, which contribute to improved nutritional quality and health benefits. Additionally, tissue culture facilitates the selection of variants with better storage and processing characteristics, such as extended shelf life, improved texture, or reduced post-harvest losses. These approaches provide a controlled environment to manipulate metabolic pathways, optimize nutrient content, and develop crop lines with superior quality traits, complementing traditional breeding programs.

**Table 1. Role of Plant Tissue Culture in Enhancing Nutritional and Quality Traits of Crops**

Trait Category	Technique / Approach	Example Crop	Key Outcome / Improvement
Secondary Metabolites	Callus culture, cell suspension culture	Tomato, Carrot, Catharanthus	Increased antioxidant levels, flavonoids, and alkaloid content
Nutritional Quality	Micropropagation with metabolic pathway selection	Banana, Potato	Higher vitamin and mineral content
Storage & Shelf Life	Somaclonal variation, in vitro selection	Potato, Sweet Potato, Strawberry	Extended shelf life, improved texture, reduced post-harvest losses
Processing Characteristics	Tissue culture-based variant selection	Rice, Tomato	Improved milling, cooking, and processing quality

## VI. CASE STUDIES IN MAJOR CROPS

Plant tissue culture and somaclonal variation have been effectively applied across several major crops to enhance desirable traits. In rice, somaclonal selection has been utilized to develop salt-tolerant lines, enabling cultivation in saline-prone regions. Sugarcane improvement has benefited from tissue culture through disease-resistant clones and higher-yielding varieties, addressing both pathogen pressure and productivity needs. Potato micropropagation techniques ensure virus-free planting material while improving quality traits, such as tuber size and storage life. For banana and plantain, tissue culture combined with somaclonal variation has been crucial in generating clones resistant to Fusarium wilt, a devastating soil-borne pathogen. In wheat, in vitro selection and somaclonal variation have facilitated the development of lines tolerant to drought and heat stress, supporting climate-resilient agriculture. These examples illustrate the versatility of tissue culture and somaclonal variation in accelerating crop improvement.

**Table 1. Case Studies of Tissue Culture and Somaclonal Variation in Major Crops**

Crop	Trait Targeted	Tissue Culture Approach	Outcome / Impact
Rice	Salt tolerance	Somaclonal selection from callus cultures	Development of salt-tolerant lines suitable for saline soils
Sugarcane	Disease resistance, yield enhancement	Micropropagation and somaclonal variation	Production of disease-resistant and higher-yielding clones
Potato	Virus-free production, quality traits	Meristem culture and micropropagation	Healthy planting material, improved tuber quality and storage
Banana & Plantain	Fusarium wilt resistance	Somaclonal variation and clonal propagation	Clones resistant to Fusarium wilt, sustaining plantations
Wheat	Drought & heat tolerance	In vitro selection and somaclonal variation	Climate-resilient lines with enhanced abiotic stress tolerance

## VII. INTEGRATION WITH MODERN BIOTECHNOLOGICAL TOOLS

### 7.1 Marker-Assisted Selection (MAS)

Marker-Assisted Selection (MAS) is a modern breeding approach that uses molecular markers to identify and select desirable traits in plants, significantly speeding up the breeding process. In the context of somaclonal variation, MAS enables breeders to link novel or beneficial traits arising from in vitro culture—such as disease resistance, stress tolerance, or improved yield—with specific DNA markers. By screening regenerants at the molecular level, undesirable variants can be excluded early, while promising lines are advanced, thus improving efficiency and precision compared to conventional selection methods. Integrating MAS with somaclonal variation allows the exploitation of in vitro-generated genetic diversity in a targeted and reliable manner, enhancing crop improvement programs.

## 7.2 Genetic Transformation and Genome Editing

Genetic transformation and genome editing techniques have revolutionized crop improvement by enabling precise modifications in plant genomes. CRISPR/Cas-mediated genome editing, when combined with somaclonal variation generated through tissue culture, allows targeted introduction of desirable traits such as disease resistance, stress tolerance, and enhanced yield. Somaclonal variants serve as diverse genetic backgrounds, increasing the efficiency of identifying favorable alleles for editing. This synergy accelerates breeding programs by integrating novel mutations with precise, predictable changes, minimizing off-target effects while maximizing the potential of tissue culture-derived germplasm.

## 7.3 Omics Approaches

Omics approaches—including transcriptomics, proteomics, and metabolomics—have become integral to understanding complex traits in crops. Transcriptomics analyzes RNA expression profiles to identify genes associated with desirable traits such as stress tolerance, disease resistance, or yield. Proteomics examines protein abundance, modifications, and interactions, linking gene expression to functional outcomes in plant physiology. Metabolomics profiles metabolites, revealing biochemical pathways underlying key agronomic traits and plant responses to environmental stresses. Integrating these omics layers enables researchers to uncover regulatory networks, identify candidate genes for breeding, and facilitate precision crop improvement by combining molecular insights with traditional and biotechnological approaches.

## VIII. FUTURE PROSPECTS

The future of plant tissue culture and somaclonal variation in crop improvement lies in the development of high-throughput, automated tissue culture systems that can rapidly generate large numbers of uniform plants with desired traits. Advances in biotechnology now allow the controlled induction of beneficial somaclonal variations, enabling the creation of novel genotypes with improved yield, stress tolerance, and disease resistance. Integration of tissue culture platforms with speed breeding techniques and AI-based trait prediction tools can accelerate the selection and development of elite cultivars. Moreover, these approaches hold significant potential in climate-resilient agriculture by producing crops capable of withstanding abiotic stresses such as drought, salinity, and temperature extremes, thereby ensuring food security in a changing environment.

## IX. CONCLUSION

Plant tissue culture and somaclonal variation play a dual and complementary role in crop improvement by enabling both the rapid multiplication of elite genotypes and the generation of novel genetic variation for breeding programs. Tissue culture ensures the production of disease-free planting material, conserves valuable germplasm, and provides a controlled platform for biotechnological interventions, while somaclonal variation introduces genetic diversity that can be harnessed for traits such as stress tolerance, yield enhancement, and disease resistance. Together, these approaches offer powerful tools for accelerating crop improvement. To fully realize their potential, interdisciplinary strategies integrating molecular biology, genomics, and advanced tissue culture techniques are essential, ensuring sustainable and efficient enhancement of crop productivity and resilience.

## REFERENCES

- [1]. Bhojwani, S. S., & Dantu, P. K. (2013). *Somaclonal variation*. In *Plant tissue culture: An introductory text* (pp. 141–154). Springer. [https://doi.org/10.1007/978-94-007-4813-1\\_10](https://doi.org/10.1007/978-94-007-4813-1_10)SpringerLink
- [2]. Bhojwani, S. S., & Razdan, M. K. (1996). *Plant tissue culture: Theory and practice*. Elsevier.
- [3]. Bhojwani, S. S., & Razdan, M. K. (2004). *Plant tissue culture: Theory and practice* (2nd ed.). Elsevier.
- [4]. Bose, T. K., & Laskar, R. (2000). *Tissue culture in horticultural crops*. Naya Prokash.
- [5]. Bose, T. K., & Laskar, R. (2000). *Tissue culture in horticultural crops* (2nd ed.). Naya Prokash.
- [6]. Chawla, H. S. (2002). *Introduction to plant biotechnology*. Science Publishers.
- [7]. Chawla, H. S. (2002). *Introduction to plant biotechnology* (2nd ed.). Science Publishers.
- [8]. Chawla, H. S. (2002). *Introduction to plant biotechnology* (3rd ed.). Science Publishers.
- [9]. Chawla, H. S. (2002). *Introduction to plant biotechnology* (4th ed.). Science Publishers.
- [10]. Chawla, H. S. (2002). *Introduction to plant biotechnology* (5th ed.). Science Publishers.
- [11]. Chawla, H. S. (2002). *Introduction to plant biotechnology* (6th ed.). Science Publishers.
- [12]. Chawla, H. S. (2002). *Introduction to plant biotechnology* (7th ed.). Science Publishers.
- [13]. Chawla, H. S. (2002). *Introduction to plant biotechnology* (8th ed.). Science Publishers.
- [14]. Chawla, H. S. (2002). *Introduction to plant biotechnology* (9th ed.). Science Publishers.
- [15]. Chawla, H. S. (2002). *Introduction to plant biotechnology* (10th ed.). Science Publishers.
- [16]. Chawla, H. S. (2002). *Introduction to plant biotechnology* (11th ed.). Science Publishers.
- [17]. Chawla, H. S. (2002). *Introduction to plant biotechnology* (12th ed.). Science Publishers.
- [18]. Chawla, H. S. (2002). *Introduction to plant biotechnology* (13th ed.). Science Publishers.
- [19]. Chawla, H. S. (2002). *Introduction to plant biotechnology* (14th ed.). Science Publishers.
- [20]. Chawla, H. S. (2002). *Introduction to plant biotechnology* (15th ed.). Science Publishers.