



Research Paper

Report of Nepal Gray Langur (*Semnopithecus schistaceus*) from Hambal area (Marmat Tehsil) of district Doda of Jammu and Kashmir.

Ajaz Ahmed Wani

Head Department of Zoology Govt. Degree College Doda, Jammu and Kashmir, India.

Abstract

Grey Langur or Hanuman Langurs the most widespread langur of Indian subcontinent are group of old world monkeys constituting entirely of genus *Semnopithecus*. These are large and fairly terrestrial inhabiting forest, open lightly woody habitat and urban areas of Indian subcontinent. Most species are found in moderate altitude, where as Nepal gray langur and Kashmir langur is found up to 4000 to 4500m. The present study is case study regarding the distribution of Nepal grey langur in erstwhile district Doda which is located in middle Himalayan chain of North West Himalaya.

Key words: Grey Langur, India, Himalaya, Jammu & Kashmir.

Received 28 Mar, 2022; Revised 06 Apr, 2022; Accepted 08 Apr, 2022 © The author(s) 2022.

Published with open access at www.questjournals.org

I. Introduction

India is a country with rich biodiversity, having 15 species and 39 sub species of non-human primates, distributed all over from Himalayan temperate forests in the north to the tropical forest in the south (Mir *et al*; 2015). Langurs are important group of mammals and are viewed as occupying important position in food chain.

Among the langurs, grey langurs are the most wide spread langurs of the Indian subcontinent. They are a group of old world monkeys constituting entirely of the genus *Semnopithecus*. All taxa had traditionally been placed in a single species *Semnopithecus entellus*. Groove 2001 recommended that several distinctive former subspecies should be given species status so that seven species are recognized. (Grove 2005) on the basis of genetic evidence suggests that Nilgiri Langur and purple faced langur which usually are placed in the genus *Trachypethicus* actually belongs to *Semnopithecus*.

The northern plain grey langur is a species of primate in the family Ercopitheidae. It is found in the lowlands north of the Godavari and Krishna rivers and Ganga. It is thought to be introduced to Western Bangladesh by Hindu Pilgrims on the bank of Jalangi river. Its natural habitat are tropical and sub tropical dry forest shrub land. It is threatened by habitat loss (Mitra's *et al*; 2008).

The Himalyan Gray Langur is distributed from central Nepal to Tibet into area of Bhutan, India and Pakistan. They are present in temperate region and live upto an elevation of 4000m, and their preferred habitat is Coniferous and broadleaf forest, but in general they show flexibility in their habitat and are even found in human dominated area. Nepal gray langur were considered a subspecies of the Northern plain gray langur. They along with seven other langur species are now considered to be separate species of same genus (Sathya Kumar *et al* 2019) used molecular study and morphological data to support the treatment of Nepal gray langur as its own species with no sub species. Nepal gray langurs are larger than most other gray langur species. They have brownish gray body with a black face. The hair framing the face is long and stark white and distant from body.

The demarcation between the head and the body is one morphological feature that makes it different from Northern Plain Langurs. Hand forelimbs tend to be light dark brown or black and bellies and chest tend to be light grey or white. They have long powerful pointed tail and fingers that are used to help move about in trees. This species of langur have tail pointed forward while Southern population tend to carry their tail pointed to the ground.

II. Observation And Discussion

The present troop of Nepal grey langur (*Semnopithecus schistaceus*), was reported from Hambal village which is located on the khellani Goha link road approximately 27 km from Doda town and is located at an altitude of and at 33°05' North to 75°24' E Longitude. . The troop comprises of 26 members which were observed around this link road at this site. This troop being mobile, is observed to move to nearby villages for feeding purpose. Grey langurs are folivores and shows variation in food consumption and food choice depending upon the habitat types and seasons. The reported troop feeds on the leaves, fruits, maize and other cereals grown in that area. Moreover there is temple at the site where this troop is present and also feed on the food items people feed them. They are shy to make it difficult to capture them in lense. They tend to run instead of walk. They are also found to run and walk intermittently.

Such species of langur was earlier reported in erstwhile district Doda, i.e Dist.Ramban by Ajaz et.al (2020) whereas Neeraj Sharma and Muzaffer (2017) reported the endangered Kashmir grey langur (*S. ajax*) in Bhaderwah of Jammu & Kashmir and reported 142 individuals in 10 troops up to 3250 m altitude are present. The Kashmir grey langur was also reported by Khaleel 2018. He studied the distribution, ecology and habitat in Dachigam National Park of J&K. This study establishes baseline information for study of distribution, habitat use and feeding ecology of *S. schistaceus* across different habitat and land uses in erstwhile district Doda.



(Nepal Grey Langoor (*SemnopethicusSchistaceus*))

Fig- *Semnopethicus schistaceus*

References

- [1]. Ajaz Ahmed wani ,Muneesh Kumar and Imteyaz Ahmed (2020) :Distribution of Nepal Gray Langur (*Semnopithecus schistaceus*) : A case study at Thokal (Shivgarh) of erstwhile district Doda of Jammu and Kashmir, IJLSci ,Vol.8(2) PP 1-3.
- [2]. Arekar KS Sathya Kumar, K Parveen Karmath (2019) Integrated taxonomy confirms the species status of Himalayan langur *Semnopithecus schistaceus*. Preprint Publication doi.10.110/60223
- [3]. Groves C (2001) Primate taxonomy. Smithsonian Institution Press.
- [4]. Groves C (2005) Mammal species of the world (Eds. D.E. Wilson and R.D Reeder) (3rd edn.), Johns Hopkins University Press. Baltimore. PP. 174-175.
- [5]. Khaleel, Mehreen (2018) Distribution, ecology and habitat use of the Himalayan grey Langur (*S. ajax*) in Jammu & Kashmir, India published in the Ruffer foundation.
- [6]. Mir ZR, Noor A, Habib B & Veeraswami GG (2015) Sesonal population density and winter survival strategies of endangered Kashmir langur (*S. ajax*) in Dachigam National Park, Kashmir, India, Springer plus 4: 562.
- [7]. Mitra S. & Molur (2008) S.e the IUCN, Red list of threatened species IUCN 2008.
- [8]. Neeraj Sharma & Muzaffer Ahmed (2017) Distribution of endangered Kashmiri Grey Langur (*Semnopithecus ajax*) in Bhaderwah Jammu and Kashmir, India Journal of Wildlife Research 2017. Vol 5. issue 1. PP.1-5.