



Research Paper

Ulam-Hyers Stability of Quadratic Functional Equation in Modular and \mathcal{M} -Spaces

Sushma Devi¹, Amrit²

¹Department of Mathematics, Kanya Mahavidyalaya Kharkhoda, Sonapat-131402, Haryana, India

²Department of Mathematics, Baba Mastnath University, Asthal Bohar-124021, Rohtak, Haryana, India

Abstract: This paper investigates the stability of new quadratic functional equations in the context of modular and \mathcal{M} -spaces. By considering the behaviour of solutions under perturbations, we establish conditions for the existence and uniqueness of stable solutions. The study extends previous results on stability theory to encompass the specific framework of quadratic functionals in modular and \mathcal{M} -spaces, providing insights into the robustness of solutions within these mathematical structures.

Keywords: \mathcal{M} -spaces, Modular space, Quadratic functional equation, Hyers-Ulam stability.

2020 AMS Classification: 39B82, 39B52, 39B72, 41A99.

I. Introduction

Ulam [15] presented the following stability problem in 1940, given metric group, number and mapping that fulfils the condition for all x, y in G , do we have an automorphism f of G and constant ϵ , relying exclusively on, such that for all x, y in G ? We refer to the automorphism equation as stable if the response is in the affirmative.

Ulam's dilemma received a positive partial solution from Hyers [5] a year later. Rassias proved a generalized version of Hyers result in 1978 in [13]. Since then, a number of authors have thoroughly examined the stability issues of several functional equations ([3–4], [6], [8], [12], [14], [16–18], [20–22]).

In fact, we also direct readers to the following papers: [1] for new developments on the conditional stability of the homomorphism equation and books; [2] for developments in Ulam's type stability; and ([7], [10]) for a comprehensive knowledge of stability theory.

The theory of modular linear spaces and its supporting theory were developed in 1950 by Nakano [11], and they were later improved upon by Koshi, Shimogaki [9], and Yamamuro [19].

In 2016, Xiuzhong Yang [22] proved the stability of a quadratic functional equation in \mathcal{M} -space. We recall some notions of \mathcal{M} -space, \mathcal{M} -normed spaces and modular space, for detailed understanding of the properties of the above spaces, the readers are required to read the book [14].

Definition 1.1. Let V be a linear space over \mathbb{F} that denotes either complex or real numbers. A non-negative valued function ϕ defined on V is called \mathcal{M} -norm (or briefly a norm) if it satisfies the following conditions:

(1) if and only if $\phi(x) = 0$.

(2) for all $x, y \in V$.

(3) $\phi(x) \geq 0$.

(4) provided $\phi(x) < \infty$.

(5) provided $\phi(x) < \infty$.

(6) provided $\phi(x) < \infty$.

A linear space equipped with \mathcal{M} -norm is called \mathcal{M} -space which will be denoted by (V, \mathcal{M}) . A complete \mathcal{M} -space is called \mathcal{M} -space.

Definition 1.2. Let V be a linear space over \mathbb{F} that denotes either complex or real numbers and \mathcal{M} . A non-negative real valued function ϕ defined on V is called \mathcal{M} -norm if it satisfies the following conditions:

(1) if and only if $\phi(x) = 0$.

(2) for all $x, y \in V$.

(3) $\phi(x) \geq 0$.

A linear space equipped with \mathcal{M} -norm is called \mathcal{M} -normed space, which will be denoted by (V, \mathcal{M}, ϕ) or briefly (V, \mathcal{M}) .

Remark 1.3. It is clear that \mathcal{M} -normed space is a special \mathcal{M} -space, and when \mathcal{M} -normed space become normed space. Comparing with the normed space, \mathcal{M} -space does not possess good metric properties, and the study of the stability

of functional equations becomes more difficult.

Definition 1.4. Let V be a vector space over $(F, \|\cdot\|)$. A generalized functional $f: V \rightarrow W$ is called a modular if for arbitrary $x, y \in V$ satisfies the following properties:

- (a) if and only if $f(x) = f(y)$
- (b) for every scalar α with $\alpha \neq 0$, $f(\alpha x) = \alpha f(x)$
- (c) , whenever $x, y \in V$ and $\alpha \in F$

If we replace (c) by (c') , whenever $x, y \in V$ and $\alpha \in F$, then, the modular f is called convex. A modular f defines a corresponding modular space, i.e., the vector space $(V, \|\cdot\|)$ given by: as .

Definition 1.5. If $(V, \|\cdot\|)$ is a modular space and the sequence $\{x_n\}$ in V , then

- (i) if $\|x_n - x_m\| \rightarrow 0$ as $n, m \rightarrow \infty$
- (ii) is known as δ -Cauchy if $\|x_n - x_m\| < \delta$ as $n, m \rightarrow \infty$
- (iii) A subset A is known as δ -complete if and only if every δ -Cauchy sequence is convergent in A .

There are many forms of the quadratic functional equation among them of great interest to us is the following:

$$(1)$$

Result 1.6. Let $(V, \|\cdot\|)$ be a linear space and a mapping f satisfies the functional equation (1), then the following results are hold:

- (1) f is linear
- (2) f is continuous

The purpose of this paper is to study the stability of equations in δ -space and Modular space. Section 2 is devoted to the study of stability of (1) in δ -space and section 3 for stability of functional equation in modular space, in more general setting such that the target spaces are δ -spaces or complete δ -normed spaces and modular space.

Throughout section 2, let $(V, \|\cdot\|)$ be a real δ -normed space and $(W, \|\cdot\|)$ be complete δ -spaces or δ -normed space. Also \mathbb{R} , and \mathbb{C} stand for the set of all real numbers, real numbers, or complex numbers and natural numbers, respectively.

II. Hyers-Ulam-Rassias Stability of (1) in δ -Space

From now on, let $(V, \|\cdot\|)$ be a real vector space and let $(W, \|\cdot\|)$ be δ -space in which there exists δ such that for all $x, y \in V$, unless we give any specific reference.

We will investigate the Hyers-Ulam-Rassias stability problem for functional equation (1). Thus, we find the condition that there exists a true quadratic function near an approximately quadratic function.

Theorem 2.1. Let $(V, \|\cdot\|)$ be a real vector space and $(W, \|\cdot\|)$ be δ -space in which there exists δ such that for all $x, y \in V$, and let f be a function such that

$$(2)$$

converges and

$$(3)$$

for all $x, y \in V$. Suppose that f satisfies

$$(4)$$

for all $x, y \in V$. Then, there exists unique quadratic function Q which satisfies (4) and the inequality

$$(5)$$

for all $x, y \in V$. Function Q is given by

$$(6)$$

for all $x, y \in V$.

Proof. Replacing x by $x+y$ in (4), we have

$$(7)$$

It follows that

$$(8)$$

for all $x, y \in V$. Replacing x by $x-y$ in (8) and by the assumption on norm, we get

$$(9)$$

Hence,

$$(10)$$

for all $x, y \in V$. Using the induction on positive integer n , we obtain that

$$(11)$$

for all $x, y \in V$. In order to prove convergence of sequence $\{Q_n(x)\}$, replace x by x/n to find that, for ,

$$(12)$$

Since the right hand side of the inequality tends to 0 as tends to infinity, sequence is a Cauchy sequence. Therefore, we may define for all . By letting in (11), we arrive at formula (5). To show that satisfies (1), replace by , respectively; then, it follows tha

$$(13)$$

Taking the limit as , we find that satisfies (1) for all . To prove the uniqueness of quadratic function subject to (5), let us assume that there exists quadratic function which satisfies (1) and inequality (5). Obviously, we have and for all and . Hence, it follows from (5) that

$$(14)$$

for all . By letting in the preceding inequality, we immediately find the uniqueness of . This completes the proof of the theorem.

Corollary 2.2. Let be a real vector space and be a complete -normed space , and let be a function such that

$$(15)$$

converges and

$$(16)$$

for all . Suppose that satisfies

$$(17)$$

for all . Then, there exists unique quadratic function which satisfies (1) and inequality

$$(18)$$

for all . Function is given by

$$(19)$$

for all .

III. Hyers-Ulam Stability of (1) in Modular Space

In this section, we take to be a real vector space and mapping to be complete modular space, we present Hyers-Ulam Stability of the functional equation (1). Let denote mapping

:

$$(20)$$

for all .

Theorem 3.1. Let be a function such that

$$(21)$$

and

$$(22)$$

for all . If a mapping with and such that

$$(23)$$

for all , then, there exists a unique quadratic mapping satisfying

$$(24)$$

for all .

Proof: Replace by in , we get

and then semi-convexity of and , we have

$$(25)$$

for all in V. So,

$$(26)$$

for all in and all non negative integers with . Thus is a Cauchy sequence in complete modular space , so there exists a as

$$(27)$$

for all in . Now

(28)

for all x, y in M , then by (27), right hand side of (28) tends to 0 as $n \rightarrow \infty$. Thus

Now,

(30)

for all integer n and for all x, y in M . Applying (29), we get the required result. Replacing x by $x+y$ in (23), we get

for all x, y .
Therefore

(31)

Taking the limit $n \rightarrow \infty$, we get

(32)

for all x, y .
To prove the uniqueness of f , let g be another quadratic mapping satisfying (24).

By taking $x=y$, we have
Therefore, the function f is unique. This completes the proof.

Corollary 3.2. If a mapping f with ϕ and ψ such that

(33)

for all x, y , then, there exists a unique quadratic mapping Q satisfying

(34)

for all x, y .

Corollary 3.3. If a mapping f with ϕ and ψ such that

(35)

for all x, y , then there exists a unique quadratic mapping Q satisfying

(36)

for all x, y .

IV. Conclusion

In this work we introduced a new quadratic functional equation of three variables and discussed about its Hyers-Ulam stability in two different spaces, M -space and Modular space respectively. In future researchers can discuss about the stability of this functional equation in other spaces.

References

- [1]. J. Brzdęk, W. Fechner, M. S. Moslehian, and J. Sikorska, "Recent developments of the conditional stability of the homomorphism equation," *Banach Journal of Mathematical Analysis*, vol. 9, no. 3, pp. 278-326, 2015.
- [2]. N. Brillouët-Belluot, J. Brzdęk, and K. Ciepliński, "On some recent developments in Ulam's type stability," *Abstract and Applied Analysis*, vol. 2012, Article ID 716936, 41 pages, 2012.
- [3]. Z. Gajda, "On stability of additive mappings," *International Journal of Mathematics and Mathematical Sciences*, vol. 14, no. 3, pp. 431-434, 1991.
- [4]. P. Găvruta, "A generalization of the Hyers-Ulam-Rassias stability of approximately additive mappings," *Journal of Mathematical Analysis and Applications*, vol. 184, no. 3, pp. 431-436, 1994.
- [5]. D. H. Hyers, "On the stability of the linear functional equation," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 27, pp. 222-224, 1941.
- [6]. S.M. Jung and J. M. Rassias, "A fixed point approach to the stability of a functional equation of the spiral of Theodorus," *Fixed Point Theory and Applications*, vol. 2008, Article ID 945010, 7 pages, 2008.
- [7]. S.M. Jung, "Hyers-Ulam-Rassias Stability of Functional Equations in Nonlinear Analysis," *Springer Optimization and Its Applications*, Springer, vol. 48, New York, NY, USA, 2011.
- [8]. S.M. Jung and P. K. Sahoo, "Hyers-Ulam stability of the quadratic equation of Pexider type," *Journal of the Korean Mathematical Society*, vol. 38, no. 3, pp. 645-656, 2001.
- [9]. S. Koshi and T. Shimogaki, "On F-norms of quasi-modular spaces," *Journal of the Faculty of Science, Hokkaido University. Series*, vol. I, no. 15, pp. 202-218, 1961.
- [10]. P. L. Kannappan, "Functional Equations and Inequalities with Applications," *Springer*, New York, NY, USA, 2009.
- [11]. H. Nakano, "Modular Semi-Ordered Linear Spaces," *Maruzen Co., Ltd.*, Tokyo, 1950.
- [12]. C. Park and J. M. Rassias, "Stability of the Jensen-type functional equation in \mathcal{A} -algebras: a fixed point approach," *Abstract and Applied Analysis*, vol. 2009, Article ID 360432, 17 pages, 2009.

- [13]. T. M. Rassias, "On the stability of the linear mapping in Banach spaces," Proceedings of the American Mathematical Society, vol. 72, no. 2, pp. 297-300, 1978.
- [14]. S. Rolewicz, "Metric Linear Spaces," Polish Scientific Publishers, Warsaw, Poland, 1972.
- [15]. S. M. Ulam, "Problems in Modern Mathematics," John Wiley & Sons, New York, NY, USA, 1960
- [16]. L. G. Wang, B. Liu, and R. Bai, "Stability of a mixed type functional equation on multi-Banach spaces: a fixed point approach," Fixed Point Theory and Applications, vol. 2010, Article ID 283827, 9 pages, 2010.
- [17]. B. Xu and J. Brzdęk, "Hyers-Ulam stability of a system of first order linear recurrences with constant coefficients," Discrete Dynamics in Nature and Society, vol. 2015, Article ID 269356, 5 pages, 2015.
- [18]. B. Xu, J. Brzdęk, and W. Zhang, "Fixed-point results and the Hyers-Ulam stability of linear equations of higher orders," Pacific Journal of Mathematics, vol. 273, no. 2, pp. 483-498, 2015.
- [19]. S. Yamamuro, "On conjugate spaces of Nakano spaces," Transactions of the American Mathematical Society, vol. 90, no. 2, pp. 291-311, 1959.
- [20]. X. Yang, L. Chang, G. Liu, and G. Shen, "Stability of functional equations in F -normed spaces," Journal of Inequalities and Applications, vol. 2015, article 112, 18 pages, 2015.
- [21]. X. Zhao, X. Yang, and C.-T. Pang, "Solution and stability of a general mixed type cubic and quartic functional equation," Journal of Function Spaces and Applications, vol. 2013, Article ID 673810, 8 pages, 2013.
- [22]. Yang, Xiuzhong. (2016), "On the Stability of Quadratic Functional Equations in F -Spaces," Journal of Function Spaces. Vol. 2016, pp 1-7. <http://dx.doi.org/10.1155/2016/5636101>.